

SURVIVAL AND GROWTH OF RESTORED PIEDMONT RIPARIAN FORESTS AS AFFECTED BY SITE PREPARATION, PLANTING STOCK, AND PLANTING AIDS

Chelsea M. Curtis, W. Michael Aust, John R. Seiler, and Brian D. Strahm¹

Abstract--Forest mitigation sites may have poor survival and growth of planted trees due to poor drainage, compacted soils, and lack of microtopography. The effects of five replications of five forestry mechanical site preparation techniques (Flat, Rip, Bed, Pit, and Mound), four regeneration sources (Direct seed, Bare root, Tubelings, and Gallon), and three planting aids (None, Mat, Tubes) on American sycamore (*Platanus occidentalis* L.) and willow oak (*Quercus phellos* L.) were examined for 2 years following establishment in order to evaluate the treatment potential for enhancing survival and growth. After 2 years, Mounding and Gallon seedlings were found to be the most beneficial treatments for American sycamore survival and growth. Bedding also proved beneficial. For willow oak, Mound and Bed were also beneficial, particularly with Bare root seedlings Gallons. The positive responses of the species to mounding and bedding were due to treatment effects on elevation on poorly drained sites, reduction of competition, and reduction of compaction.

INTRODUCTION

The Federal Water Pollution Control Act of 1972 and subsequent amendments and interpretations have resulted in policies which require wetland restoration or creation to offset wetland losses caused by activities such as urbanization (Stolt and others 2000). Wetland creation projects have a relatively poor track record for success, thus it is common to have wetland mitigation ratios of 2:1 or 3:1 (Brown and Lant 1999). The relatively poor success rates are caused by a variety of problems, including: poor recognition of site conditions which results in poor species selection; sites with compacted soil conditions, which inhibit soil water movement and root penetration; excessively wet sites that may kill or suppress growth of desired tree species; and lack of topography which may limit the survival and growth of planted tree species (Bailey and others 2007).

Forest managers have been facing similar regeneration problems on such sites. Harvested sites are commonly compacted and poorly drained, yet silviculturalists have overcome these limitations with a variety of mechanical site preparation techniques (Aust and others 1998). For example, both mounding and bedding have been widely used across the eastern United States since the 1950s and 1960s to overcome lack of relief and soil compaction on wet sites (Lof and others 2012, Miwa and others 2004). Similarly, riparian restoration efforts often have

site limitations that are overcome by using alternative planting sources or planting aids. Interestingly, there has been little technical transfer between forest managers and the wetland restoration community.

The literature indicates that wetland restoration efforts could be enhanced with increased use of silvicultural tools. Therefore, the objective of this research project is to quantify the effects of mechanical site preparation, regeneration source, and planting aids on the survival and growth of two species commonly used on mitigation sites: the early successional species American sycamore (*Platanus occidentalis* L.) and the later successional species willow oak (*Quercus phellos* L.).

MATERIALS AND METHODS

Study Site

The study site is located in the Piedmont physiographic province on the Virginia Tech R.J. Reynolds Homestead Forest Research Extension Center near Critz, VA. Much of this 280-ha area was converted to tobacco plantations during the 1800s, and the specific riparian area was subjected to agriculture and excessive compaction by a recent soil compaction research project. Thus the area has compacted soils, lack of relief, and is excessively wet during winter and spring. Soil series in the study include Augusta (fine-loamy, mixed semiactive, thermic Aeric Endoaquults), French (fine loamy over sandy, mixed, active

¹Graduate Research Assistant, Professor, Professor, and Assistant Professor, respectively, Virginia Polytechnic Institute and State University, Department of Forest Resources and Environmental Conservation, Blacksburg, VA 24061.

mesic Fluvaquentic Dystrudepts), and Roanoke (fine, mixed, semiaactive thermic Typic Endoaquults). The site is located in the floodplain of a first-order perennial stream with a 500-ha watershed. Flooding to a depth of approximately 25 cm occurs during most spring seasons.

Treatments

Five site preparation treatments were established (Flat, Rip, Bed, Pit, Mound). The Flat treatment consisted of surface tillage with a disk harrow to reduce herbaceous competition. The Rip treatment consisted of subsoiling with a 30-cm ripping shank underneath the planting zones. The Bed treatments were made with the blade on a bulldozer. The Pit and Mound treatments were created in the same area: a tractor-mounted backhoe was used to excavate pit material (approximately 40 cm) to create an adjacent mound of approximate 40 cm. Four regeneration sources (Seed, Bare root, Tubeling, Gallon) were superimposed across all site preparation treatments. For the Seed treatment, three acorns or a finger pinch of American sycamore seeds were planted. The seeds were collected from piedmont seed sources approximately 3 months prior to planting. The bare-root seedlings were purchased from commercial nurseries and planted with dibble bars. Tubeling and Gallon containers were purchased from a commercial nursery and planted with spades. Three levels of planting aids were applied to all combinations of site preparation and regeneration sources (None, Tubes, and Mats). Tubes consisted of 1-m planting tubes, and mats were 50- by 50-cm geotextile fabric. Seeds and seedlings were planted in May 2011, and planting aids were installed in June 2012. Minimal herbaceous control was conducted during summer 2011 and 2012.

Survival and Growth Parameters

Survival and growth indices were measured after one and two growing seasons (2011, 2012) simply recording if the individual had survived. Growth parameters included average ground line diameter based on caliper measurements from two directions and total tree height

measured to the nearest 1 cm with a height poles. For trees taller than 1.3 m, diameter at breast height (d.b.h., cm) was also recorded. The diameter and height measures were subsequently converted to a biomass index in cm^3 based on d^2h geometry.

Statistical Design and Analysis

The study is arranged as a split-split plot within a Randomized Complete Block Design. Five blocks were established for each of the two species. The main effects are five site preparation treatments (Flat, Rip, Bed, Pit, and Mound). The Pit and Mound treatments were established together but were analyzed as two treatments. The split plot was comprised of the four regeneration sources. The second split consisted of the three planting aids. For each experimental unit, four units (seed or seedling) were established. Thus, for both American sycamore and willow oak, approximately 1,200 trees or seeds were planted (5 blocks x 5 site preparation treatments x 4 regeneration sources x 3 planting aids x 4 trees per seeds = 1,200). Survival and growth parameters were analyzed via analysis of variance (ANOVA) and statistically different means were separated with a Tukeys HSD test.

RESULTS AND DISCUSSION

American sycamore survival was consistently between 62 and 66 percent across the site preparation treatments during years 1 and 2 with the exception of the Pit treatment, which had significantly lower survival (table 1). American sycamore's biomass index was significantly reduced by the Pit treatment during both years and was significantly increased by the Mound site preparation treatment (table 1). Not unexpectedly, American sycamore survival percentages were favored by the Tubeling and Gallon regeneration sources and were very low for direct seeding (table 2). Biomass values followed the same general trends: lowest for Direct seeding, followed by Bare root and Tubeling, and greatest in the Gallon regeneration sources (table 2). Examination of the effects of planting aids on American sycamore survival and growth indicated that the

Table 1--Effects of site preparation on survival and biomass indices for American sycamore on a Piedmont forest restoration site during years 1 and 2. Mean values within a column followed by a different letter are significantly different at $\alpha \leq 0.10$

| Site preparation treatment | -----Survival----- | | ---Biomass index--- | |
|----------------------------|--------------------|---------------|----------------------------|---------------|
| | Year 1 | Year 2 | Year 1 | Year 2 |
| | -----percent----- | | -----cm ³ ----- | |
| Flat | 68b | 68b | 496b | 1487b |
| Rip | 68b | 68b | 549b | 1900b |
| Bed | 66b | 66b | 649b | 2497b |
| Pit | 62a | 59a | 333a | 1232a |
| Mound | 68b | 68b | 913c | 3811c |
| p | ≤ 0.0001 | ≤ 0.0561 | ≤ 0.0001 | ≤ 0.0001 |

Table 2--Effects of regeneration sources on survival and biomass indices for American sycamore on a piedmont forest restoration site during years 1 and 2. Mean values within a column followed by a different letter are significantly different at $\alpha \leq 0.10$

| Regeneration source treatment | -----Survival----- | | ---Biomass index--- | |
|-------------------------------|--------------------|---------------|----------------------------|---------------|
| | Year 1 | Year 2 | Year 1 | Year 2 |
| | -----percent----- | | -----cm ³ ----- | |
| Direct seed | 36a | 16a | 10a | 10a |
| Bare root | 72b | 58b | 196a | 933b |
| Tubling | 85c | 85c | 445b | 1896c |
| Gallon | 88c | 88c | 1023c | 3728d |
| p | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.0001 |

planting Mats and Tubes slightly increase survival, and the Mat increased biomass (table 3).

Willow oak survival was also reduced by the Pit treatment (table 4). Mound treatment positively increased biomass for willow oak during year 1. During year 2, the pattern continued, but the differences were not significant. Bare root and Gallon both provided good survival for willow oak (table 5), and the Direct seeding survivals were lower. However, survival of the relatively larger-seeded willow oak was better with direct seeding than for American sycamore. During year 1, the gallon regeneration source offered the best biomass growth for willow oak, but by year 2, regeneration sources effects on willow oak biomass were not significant (table 5). The Mat planting aid provided both enhanced survival and biomass for willow oak (table 6).

The efficacy of any given treatment will be due to a combination of the treatment effects on both survival and growth. Therefore, we created a unitless performance index which is the product of survival and biomass for both American sycamore (table 7) and willow oak (table 8). We then examined the top 25 percent of all treatments to select the "best" treatments. For American sycamore, the Mound treatment followed by the Bed site preparation treatment combined with large Gallon containerized seedlings clearly performed the best. There was no clear pattern with planting aids for American sycamore. For willow oak the Mound and Bed treatments worked well for the Gallon containers, but the Bare root seedlings also performed well. For both species, the Mound and Bed treatments elevated the seedlings in the poorly drained soils and favored tree survival. The Mound and Bed treatments also alleviated soil compaction. The Mound treatment

Table 3--Effects of planting aids on survival, diameters, heights, and biomass indices for American sycamore on a piedmont forest restoration site during years 1 and 2. Mean values within a column followed by a different letter are significantly different at $\alpha \leq 0.10$. Year 1 biomass indices were not significantly different

| Planting aid treatment | -----Survival----- | | --Biomass index-- | |
|------------------------|--------------------|---------------|----------------------------|---------------|
| | Year 1 | Year 2 | Year 1 | Year 2 |
| | -----percent----- | | -----cm ³ ----- | |
| None | 71a | 63a | 600 | 2230ab |
| Tube | 76b | 66ab | 533 | 1694a |
| Mat | 77b | 70b | 649 | 2632b |
| p | ≤ 0.006 | ≤ 0.0001 | ≤ 0.6370 | ≤ 0.0144 |

Table 4--Effects of site preparation on survival and biomass indices for willow oak on a piedmont forest restoration site during years 1 and 2. Mean values within a column followed by a different letter are significantly different at $\alpha \leq 0.10$. Year 2 biomass indices were not significantly different

| Site preparation treatment | -----Survival----- | | ---Biomass index--- | |
|----------------------------|--------------------|---------------|----------------------------|---------------|
| | Year 1 | Year 2 | Year 1 | Year 2 |
| | -----percent----- | | -----cm ³ ----- | |
| Flat | 77b | 56b | 53a | 2193 |
| Rip | 79b | 60b | 67b | 1471 |
| Bed | 80b | 80b | 69b | 1435 |
| Pit | 67a | 67a | 70b | 440 |
| Mound | 83b | 83b | 84c | 2001 |
| p | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.1507 |

Table 5--Effects of regeneration sources on survival and biomass indices for willow oak on a piedmont forest restoration site during years 1 and 2. Mean values within a column followed by a different letter are significantly different at $\alpha \leq 0.10$. Year 2 biomass indices were not significantly different

| Regeneration source treatment | -----Survival----- | | ---Biomass index--- | |
|-------------------------------|--------------------|---------------|----------------------------|---------------|
| | Year 1 | Year 2 | Year 1 | Year 2 |
| | -----percent----- | | -----cm ³ ----- | |
| Direct seed | 61a | 31a | 0.5a | 1240 |
| Bare root | 86c | 86c | 65c | 2278 |
| Tubling | 74b | 65b | 5b | 1435 |
| Gallon | 84c | 84c | 127d | 1104 |
| p | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.2038 |

Table 6--Effects of planting aids on survival, diameters, heights, and biomass indices for willow oak on a piedmont forest restoration site during years 1 and 2. Mean values within a column followed by a different letter are significantly different at $\alpha \leq 0.10$. Year 1 biomass indices were not significantly different

| Planting aid treatment | -----Survival----- | | --Biomass index-- | |
|------------------------|--------------------|---------------|----------------------------|---------------|
| | Year 1 | Year 2 | Year 1 | Year 2 |
| | -----percent----- | | -----cm ³ ----- | |
| None | 74a | 51a | 67 | 718a |
| Tube | 78ab | 60ab | 67 | 1620b |
| Mat | 82b | 69b | 67 | 2154c |
| p | ≤ 0.0001 | ≤ 0.0001 | ≤ 0.3323 | ≤ 0.0011 |

Table 7--American sycamore performance index (biomass x survival) at 2 years. Numbers with asterisk represent the top 25 percent of all treatment combinations for performance

| Regeneration source | Planting aid | -----Site preparation treatment----- | | | | |
|---------------------|--------------|--------------------------------------|-------|-------|-------|-------|
| | | Flat | Rip | Bed | Pit | Mound |
| Direct seed | None | 1 | 156 | 17 | 1 | 337 |
| | Tube | 1 | 438 | 112 | 76 | 76 |
| | Mat | 2 | 3 | 25 | 7 | 11 |
| Bare root | None | 557 | 550 | 670 | 138 | 2023* |
| | Tube | 426 | 530 | 770 | 257 | 1370 |
| | Mat | 402 | 451 | 250 | 92 | 852 |
| Tubling | None | 645 | 1523 | 2238* | 382 | 2234* |
| | Tube | 721 | 831 | 1084 | 443 | 874 |
| | Mat | 893 | 1616 | 1799 | 875 | 3119* |
| Gallon | None | 2193* | 2208* | 1923* | 1803 | 3113* |
| | Tube | 1684 | 2038* | 2735* | 1393 | 3456* |
| | Mat | 1592 | 1905* | 3532* | 2042* | 6234* |

Table 8--Willow oak performance index (biomass x survival) at 2 years. Numbers with asterisk represent the top 25 percent of all treatment combinations for performance

| Regeneration source | Planting aid | -----Site preparation treatment----- | | | | |
|---------------------|--------------|--------------------------------------|-------|-------|-----|-------|
| | | Flat | Rip | Bed | Pit | Mound |
| Direct seed | None | 111 | 4 | 4 | 1 | 5 |
| | Tube | 20 | 86 | 10 | 1 | 31 |
| | Mat | 1 | 10 | 15 | 1 | 39 |
| Bare root | None | 398 | 748 | 1541* | 145 | 1624* |
| | Tube | 2173* | 516 | 1015* | 76 | 1025* |
| | Mat | 669 | 674 | 787 | 237 | 1424* |
| Tubling | None | 52 | 127 | 33 | 1 | 153 |
| | Tube | 101 | 108 | 38 | 45 | 118 |
| | Mat | 116 | 53 | 13 | 8 | 390 |
| Gallon | None | 727 | 1067* | 987* | 287 | 1480* |
| | Tube | 676 | 985* | 1157* | 518 | 1354* |
| | Mat | 888 | 972* | 1168* | 446 | 1201* |

provided some competition control by burying the seeds of competitors deeper than they could survive. Currently, the use of large seedlings is common practice on mitigation sites, but these data indicate that Mound and Bed also offer significant potential for improving wetland mitigation. The wetland mitigation community typically has close working ties with equipment contractors, thus locating excavator operators should be relatively straightforward. The use of Mounds offers significant potential to overcome the typical problems encountered on Piedmont mitigation sites.

ACKNOWLEDGEMENTS

This research was funded by the Peterson Family Foundation, Wetlands Studies and Solutions, Incorporated, the R.J. Reynolds Forest Research Extension Center, and Virginia Tech.

LITERATURE CITED

- Aust, W.M.; Burger, J.A.; McKee, W.H., Jr. [and others]. 1998. Bedding and fertilization ameliorate effects of designated wet-weather skid trails after four years for loblolly pine (*Pinus taeda*) plantations. *Southern Journal of Applied Forestry*. 22(4): 222-226.
- Bailey, D.E.; Perry, J.E.; Daniels, W.L. 2007. Vegetation dynamics in response to organic matter loading rates in a created freshwater wetland in southeastern Virginia. *Wetlands*. 27(4): 936-950.
- Brown, P.H.; Lant, C.L. 1999. The effect of wetland mitigation banking on the achievement of no-net-loss. *Environmental Management*. 23(3): 333-345.
- Lof, M.; Dey, D.C.; Navarro, R.M.; Jacobs, D.F. 2012. Mechanical site preparation for forest restoration. *New Forests*. 48: 825-848.
- Miwa, M.; Aust, W.M.; Burger, J.A. [and others]. 2004. Wet-weather timber harvesting and site preparation effects on coastal plain sites: a review. *Southern Journal of Applied Forestry*. 28: 137-151.
- Stolt, M.H.; Genthner, M.H.; Daniels, W.L. [and others]. 2000. Comparison of soil and other environmental conditions in constructed and adjacent palustrine reference wetlands. *Wetlands*. 20(4): 671-683.